Abstract Submitted for the <u>26th Annual DAMOP Meeting</u> of the American Physical Society May 16-19, 1995

Suggested title of session in which paper should be placed General Theory of Electronic Structure

Relativistic Configuration Interaction Calculations of the 2s - $2p_{3/2}$ Transitions in Highly-Ionized Uranium.* K. T. CHENG, M. H. CHEN, Lawrence Livermore National Lab., W. R. JOHNSON, U. of Notre Dame. — A large-scale, relativistic configuration interaction (CI) method with B-spline basis sets is used to calculate the 2s - $2p_{3/2}$ transition energies for highly-ionized Li-like, Be-like, B-like, and C-like uranium. These CI calculations are based on the relativistic no-pair Hamiltonian and include contributions from the Coulomb and the retarded Breit interactions. Quantum electrodynamic corrections are also calculated nonperturbatively in external Dirac-Slater potentials. Our results are in very good agreement with recent high precision experimental data taken at the SuperEBIT facility.

*Supported in part under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract No. W-7405-ENG-48 and by the National Science Foundation Grant No. 92-04089.

(x) Prefer Poster Session

Submitted by

Kwok-Tsang Cheng Lawrence Livermore National Lab. P. O. Box 808, L-59 Livermore, CA 94550 Phone: (510) 423-8659 FAX: (510) 422-5102 E-mail: ktcheng@llnl.gov